

References

- Aarts, P. B., Jongerius, P. H., Geerdink, Y. A., van Limbeek, J., & Geurts, A. C. (2010). [Effectiveness of modified constraint-induced movement therapy in children with unilateral spastic cerebral palsy: A randomized controlled trial.](#) *Neurorehabilitation and Neural Repair*, 24, 509-518.
- Blakeslee, S. (2004, February 3). [Regimen benefits children with cerebral palsy.](#) *The New York Times*.
- Brady, K., & Garcia, T. (2009). [Constraint-induced movement therapy \(CIMT\): Pediatric applications.](#) *Developmental Disabilities Research Reviews*, 15, 102-111.
- Case-Smith, J., DeLuca, S.C., Stevenson, R., & Ramey, S. L. (2012). [Multicenter randomized controlled trial of pediatric constraint-induced movement therapy: 6- month follow-up.](#) *American Journal of Occupational Therapy*, 66, 15-23.
- Charles, J., Lavinder, G., & Gordon, A. M. (2001). [Effects of constraint-induced therapy on hand function in children with hemiplegic cerebral palsy.](#) *Pediatric Physical Therapy*, 13(2), 68-76.
- Cope, S. M., Forst, H. C., Bibis, D., & Liu, X.-C. (2008). [Modified constraint-induced movement therapy for a 12-month-old child with hemiplegia: A case report.](#) *American Journal of Occupational Therapy*, 62, 430-437.
- DeLuca, S. C., Echols, K., Ramey, S. L. & Taub, E. (2003). [Pediatric constraint-induced movement therapy for a young child with cerebral palsy: Two episodes of care.](#) *Physical Therapy*, 83(11), 1003-1014.
- Eliasson, A. C., Gordon, A. M., & Forssberg, H. (2000). [Impaired force coordination during object release in children with hemiplegic cerebral palsy.](#) *Developmental Medicine and*

Child Neurology, 42, 228-234.

- Facchin, P., Rosa-Rizzotto, M., Turconi, A.C., Pagliano, E., Fazzi, E., Stortini, M., & Fedrizzi, E. (2009). [Multisite trial on efficacy of constraint-induced movement therapy in children with hemiplegia: Study design and methodology](#). *American Journal of Physical Medicine & Rehabilitation*, 88(3), 216-230.
- Gilmore, R., Ziviani, J., Sakzewski, L., Shields, N., & Boyd, R. (2010). [A balancing act: Children's experience of modified constraint-induced movement therapy](#). *Developmental Neurorehabilitation*, 13(2), 88-94.
- Gordon, A. M., Charles, J., & Wolf, S. L. (2005). [Methods of constraint-induced movement therapy for children with hemiplegic cerebral palsy: Development of a child-friendly intervention for improving upper-extremity function](#). *Archives of Physical Medicine and Rehabilitation*, 86(4), 837-844.
- Huang, H.-h., Fethers, L., Hale, J., & McBride, A. (2009). [Bound for Success: A Systematic Review of Constraint-Induced Movement Therapy in Children With Cerebral Palsy Supports Improved Arm and Hand Use](#). *Physical Therapy*, 89(11), 1126-1141.
- Knis-Matthews, L., Falzarano, M., Baum, D., Manganiello, J., Patel, S., & Winters, L. (2011). [Parents' experiences with services and treatment for their children diagnosed with cerebral palsy](#). *Physical and Occupational Therapy in Pediatrics*, 31(3), 263-274.
- Martin, A., Burtner, P. A., Poole, J., & Phillips, J. (2008). [Case report: ICF-level changes in a preschooler after constraint-induced movement therapy](#). *American Journal of Occupational Therapy*, 62(3), 282-288. doi: 10.5014/ajot.62.3.282
- Ostendorf, C.G. & Wolf, S. L. (1981). [Effect of forced use of the upper extremity of a hemiplegic patient on changes in function](#). *Physical Therapy*, 61, 1022-1028.

- Page, S.J. (2004). [Modified constraint-induced therapy for hemiparesis: A review](#). *Critical Reviews in Physical and Rehabilitation Medicine*, 16(1), 31-38.
- Plummer, A. C. (2003). [Constraint-induced therapy and the motor learning literature that underpins its application](#). *Physical Therapy Reviews*, 8, 143-149.
- Ramachandran, S. & Thakur, P. (2011). [Upper extremity constraint-induced movement therapy in infantile hemiplegia](#). *Journal of Pediatric Neurosciences*, 6(1) 29-31.
- Ries, J. D. & Leonard, R. (2006). [Is there evidence to support the use of constraint- induced therapy to improve the quality or quantity of upper extremity function of a 2 ½- year-old girl with congenital hemiparesis? If so what are the optimal parameters of this intervention?](#) *Physical Therapy Journal*, 86(5) 746-752.
- Sakzewski, L., Ziviani, J., Abbott, D. F., Macdonell, R. A., Jackson, G. D., & Boyd, R. N. (2011a). [Equivalent retention of gains at 1 year after training with constraint-induced or bimanual therapy in children with unilateral cerebral palsy](#). *Neurorehabilitation and Neural Repair*, 25(7), 664-671. doi: 10.1177/1545968311400093
- Sakzewski, L., Ziviani, J., Abbott, D. F., Macdonell, R. A., Jackson, G. D., & Boyd, R. N. (2011b). [Participation Outcomes in a Randomized Trial of 2 Models of Upper-Limb Rehabilitation for Children With Congenital Hemiplegia](#). *Archives of Physical Medicine & Rehabilitation*, 92(4), 531-539. doi: 10.1016/j.apmr.2010.11.022
- Sakzewski, L., Ziviani, J., & Boyd, R. N. (2011c). [Best responders after intensive upper-limb training for children with unilateral cerebral palsy](#). *Archives of Physical Medicine & Rehabilitation*, 92(4), 578-584. doi: 10.1016/j.apmr.2010.12.003
- Sakzewski, L., Carlon, S., Shields, N., Ziviani, J., & Ware, R. S. (2012). [Impact of intensive upper limb rehabilitation on quality of life: A randomized trial in children with unilateral](#)

- [cerebral palsy](#). *Developmental medicine and child neurology*, 54(5), 415-423.
- Stearns, G., Burtner, P., Keenan, K., Qualls, C., & Phillips, J. (2009). [Effects of constraint-induced movement therapy on hand skills and muscle recruitment of children with spastic hemiplegic cerebral palsy](#). *Neurorehabilitation*, 24(2), 95-108. doi:10.3233/NRE-2009-0459
- Taub, E. (2004). [Efficacy of constraint-induced movement therapy for children with cerebral palsy with asymmetric motor impairment](#). *Pediatrics*, 113(2), 305-312.
- Taub, E., Griffin, A., Uswatte, G., Gammons, K., Nick, J., & Law, C. (2011). [Treatment of congenital hemiparesis with pediatric constraint-induced movement therapy](#). *Journal of Child Neurology*, 26(9), 1163-1173.
- Taub, E. & Uswatte, G. (1999). [Constraint-induced movement therapy: A new family of techniques with broad application to physical rehabilitation -- a clinical review](#). *Journal of Rehabilitation Research and Development*, 36(3), 25-32.
- Taub, E., & Wolf, S. L., (1997). [Constraint induced movement techniques to facilitate upper extremity use in stroke patients](#). *Topics in Stroke Rehabilitation*, 3, 38-61.
- Wallen, M., Ziviani, J., Naylor, O., Evans, R., Novak, I., & Herbert, R. D. (2011). [Modified constraint-induced therapy for children with hemiplegic cerebral palsy: A randomized trial](#). *Developmental Medicine & Child Neurology*, 53(12), 1091-1099.
doi:10.1111/j.1469-8749.2011.04086.x
- Willis, J. K., Morello, A., Davie, A., Rice, J. C. & Bennett, J. T. (2002). [Forced use treatment of childhood hemiparesis](#). *Pediatrics*, 110(1), 94-96.
- Wolf, S. L., LeCraw, D. E., Barton, L. A., & Jann, B. B. (1989). [Forced use of hemiplegic upper extremities to reverse the effect of learned nonuse among chronic stroke and head-injured](#)

[patients](#). *Experimental Neurology*, 104, 125-132.

Wu, W., Hung, J., Tseng, C., & Huang, Y. (2013). [Group constraint-induced movement therapy for children with hemiplegic cerebral palsy: A pilot study](#). *American Journal of Occupational Therapy*, 67(2), 201-208. doi: 10.5014/ajot.2013.004374